



TITLE:

On Gauss maps in positive characteristics

AUTHOR(S):

Ito, Atsushi

CITATION:

Ito, Atsushi. On Gauss maps in positive characteristics. 代数幾何学シンポジウム記録 2014, 2014: 99-106

ISSUE DATE:

2014

URL:

<http://hdl.handle.net/2433/215016>

RIGHT:

ON GAUSS MAPS IN POSITIVE CHARACTERISTICS

ATSUSHI ITO

ABSTRACT. In [FI2], K. Furukawa and the author study Gauss maps in positive characteristics. In this note, we review [FI2].

1. INTRODUCTION

Let $X \subset \mathbb{P}^N$ be an n -dimensional projective variety over an algebraically closed field of characteristic $p \geq 0$. Then we can define the *Gauss map* γ_X of X to be the rational map

$$\gamma_X : X \dashrightarrow \mathbb{G}(n, \mathbb{P}^N)$$

which maps a smooth point $x \in X$ to the embedded tangent space $\mathbb{T}_x X \subset \mathbb{P}^N$. Here $\mathbb{G}(n, \mathbb{P}^N)$ is the Grassmannian which parametrizes n -dimensional linear subvarieties of \mathbb{P}^N .

Example 1.1. (1) If $X \subset \mathbb{P}^N$ is an n -dimensional linear subvariety, $\mathbb{T}_x X = X$ holds for any $x \in X$, i.e. γ_X is a constant map. In fact, γ_X is a constant map for a variety $X \subset \mathbb{P}^N$ if and only if X is linear.
(2) Let $X \subset \mathbb{P}^N$ be the projective cone of $Y \subset \mathbb{P}^{N-1}$ with the vertex $v \in \mathbb{P}^N$. Then γ_X is constant on the line \overline{vy} for each $y \in Y$.
(3) If $C \subset \mathbb{P}^N$ is a nonlinear curve in $p = 0$, it is known that γ_C is birational onto the image. Geometrically, this means that a general tangent line of C is tangent to C at only one point.

If we take a local parametrization of X , we can compute γ_X explicitly as follows.

Example 1.2. Let $X := \mathbb{P}^1 \hookrightarrow \mathbb{P}^N$ be the twisted cubic curve, which is locally parametrized as $[1 : t] \mapsto [1 : t : t^2 : t^3] \in \mathbb{P}^3$. For $x = [1 : t]$, the embedded tangent space $\mathbb{T}_x X \subset \mathbb{P}^3$ is the line spanned by $[1 : t : t^2 : t^3]$ and $[0 : 1 : 2t : 3t^2]$, where the latter point is obtained by differentiating $1, t, t^2, t^3$ by t . Then we have

$$\begin{bmatrix} 1 & t & t^2 & t^3 \\ 0 & 1 & 2t & 3t^2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -t^2 & -2t^3 \\ 0 & 1 & 2t & 3t^2 \end{bmatrix},$$

where \sim means that the former matrix is transformed to the latter one by elementary transformations of rows. Hence the Gauss map $\gamma_X : X \rightarrow$

$\mathbb{G}(1, \mathbb{P}^3)$ is locally described as

$$[1 : t] \mapsto \begin{bmatrix} -t^2 & -2t^3 \\ 2t & 3t^2 \end{bmatrix} \in \mathbb{A}^4 \subset \mathbb{G}(1, \mathbb{P}^3).$$

By this description, γ_X is an embedding if $p \neq 2$, but is not birational if $p = 2$.

Example 1.3. Assume $p > 0$ and let c be a positive integer prime to p . Let $X := \mathbb{P}^1 \hookrightarrow \mathbb{P}^3$, which is parametrized as $[1 : t] \mapsto [1 : t : t^{cp} : t^{cp+1}]$. Then $\gamma_X : X \rightarrow \mathbb{G}(1, \mathbb{P}^3)$ is locally described as

$$[1 : t] \mapsto \begin{bmatrix} t^{cp} & 0 \\ 0 & t^{cp} \end{bmatrix} \in \mathbb{A}^4 \subset \mathbb{G}(1, \mathbb{P}^3).$$

Thus a general fiber of γ_X is (non-reduced) c points. Geometrically, this means that a general tangent line of X is tangent to X at c points.

Remark 1.4. In [FI1], Furukawa and the author study Gauss maps of toric varieties.

About general fibers of Gauss maps, the following are known: If $p = 0$, (the closure of) a general fiber of γ_X is a linear subvariety for any X (see P. Griffiths and J. Harris [GH, (2.10)], F.L. Zak [Za, I, 2.3. Theorem (c)]). We will omit the words “the closure of” sometimes for simplicity. In particular, X is birational to $\gamma_X(X) \times \mathbb{A}^\delta$ for $\delta = \dim X - \dim \gamma_X(X)$, and the field extension $K(X)/K(\gamma_X(X))$ between function fields is purely transcendental. In $p > 0$, Furukawa [Fur] show that the same statement holds if γ_X is separable.

On the other hand, a general fiber of Gauss maps can be a union of points as we see in Example 1.3 (such examples were first found by H. Kaji [Ka1, Example 4.1] and J. Rathmann [Ra, Example 2.13]), and can be a nonlinear variety (S. Fukasawa [Fuk1, Section 7]) in $p > 0$. More generally, Fukasawa show that any projective variety appears as a general fiber of γ_X for some X :

Theorem 1.5 ([Fuk2]). *Assume $p > 0$. Let $F \subset \mathbb{P}^{N'}$ be a projective variety. If $n \geq N'$, there exists an n -dimensional projective variety $X \subset \mathbb{P}^N$ such that a general fiber of γ_X (with the reduced structure) is projectively equivalent to $F \subset \mathbb{P}^{N'}$.*

The following is an example of γ_X whose general fiber is a conic.

Example 1.6. Let $\varphi : (k^\times)^2 \hookrightarrow \mathbb{P}^3$ be an embedding defined by $(s, t) \mapsto [1 : s : s^2t : s^2t^{p+1}]$ for $p > 0$. Set $X = \overline{\varphi((k^\times)^2)} \subset \mathbb{P}^3$. Then $\gamma_X : X \dashrightarrow \mathbb{G}(2, \mathbb{P}^3) \simeq \mathbb{P}^3$ is locally described as

$$(s, t) \mapsto (0, 0, t^p) \in \mathbb{A}^3 \subset \mathbb{G}(2, \mathbb{P}^3).$$

Hence the fiber of γ_X over $(0, 0, a) \in \mathbb{A}^3$ for $a \neq 0$ is

$$\varphi(k^\times \times \{a^{1/p}\}) = \{[1 : s : a^{1/p}s^2 : a^{p+1/p}s^2] \mid s \in k^\times\} \subset \mathbb{P}^3,$$

which is a conic.

By these results, we see that fibers of Gauss maps in $p > 0$ are quite different from those in $p = 0$. Hence we ask the following question.

Question 1.7. *What kind of difference appears about Gauss maps in $p > 0$ and $p = 0$?*

In [FI2], Furukawa and the author study Question 1.7 from the view point of images $\gamma_X(X)$, general fibers, and field extensions $K(X)/K(\gamma_X(X))$ induced by γ_X . In this article, we explain the results in [FI2].

Acknowledgments. The author would like to express his gratitude to organizers Professors Keiji Oguiso, Hisanori Ohashi, and Kentaro Mitsui for giving me the opportunity to talk in this memorial conference. The author is supported by the Grant-in-Aid for JSPS fellows No. 26-1881.

2. MAIN RESULTS

I.M. Landsberg and J. Piontkowski independently gave a necessary and sufficient condition for a subvariety $Y \subset \mathbb{G}(n, \mathbb{P}^N)$ to be the image of some n -dimensional variety $X \subset \mathbb{P}^N$ in $p = 0$. In other word, they characterized subvarieties of $\mathbb{G}(n, \mathbb{P}^N)$ which are written as the image of some Gauss map (see [FP, 2.4.7] and [IL, Theorem 3.4.8]). By their characterization, we know that in $p = 0$,

- (1) not all subvariety of $\mathbb{G}(n, \mathbb{P}^N)$ is the image of some gauss maps. That is, there exists a subvariety $Y \subset \mathbb{G}(n, \mathbb{P}^N)$ with $\dim Y \leq n$ such that $Y \neq \overline{\gamma_X(X)}$ for any n -dimensional variety $X \subset \mathbb{P}^N$.
- (2) If $Y = \overline{\gamma_X(X)}$ holds for some X , such X is uniquely determined by Y .

Furukawa [Fur] generalized their characterization to separable Gauss maps in any characteristics. In particular, (1), (2) hold under the assumption that γ_X is separable in $p \geq 0$.

On the other hand, Kaji show the following proposition about images of Gauss maps of curves in $p > 0$:

Proposition 2.1 ([Ka2, Appendix]). *Assume $p > 0$. For any curve $Y \subset \mathbb{G}(1, \mathbb{P}^N)$, there exists a curve $X \subset \mathbb{P}^N$ such that $\overline{\gamma_X(X)} = Y$.*

More precisely, for any curve $Y \subset \mathbb{G}(1, \mathbb{P}^N)$ and any finite and inseparable field extension $L/K(Y)$, there exists a curve $X \subset \mathbb{P}^N$ such that $\overline{\gamma_X(X)} = Y$, $K(X) = L$, and the field extension $K(X)/K(Y)$ induced by γ_X coincides with the given extension $L/K(Y)$.

Remark 2.2. For a fixed curve $Y \subset \mathbb{G}(1, \mathbb{P}^N)$, there exist infinitely many field extensions $L/K(Y)$ which are finite and inseparable. Thus there are infinitely many curves $X \subset \mathbb{P}^N$ such that $\overline{\gamma_X(X)} = Y$ by Proposition 2.1. Hence the above (1), (2) does not hold in $p > 0$.

We generalize Proposition 2.1 to higher dimensional X, Y . In the rest of this article, a field means a finitely generated field over k . For a field extension L/K , we denote by $\delta_{L/K}$ the natural L -linear map $\Omega_{K/k} \otimes_K L \rightarrow \Omega_{L/k}$. The following is our main result.

Theorem 2.3. *Assume $p > 0$. For any variety $Y \subset \mathbb{G}(1, \mathbb{P}^N)$ with $1 \leq \dim Y \leq n$, there exists an n -dimensional variety $X \subset \mathbb{P}^N$ such that $\overline{\gamma_X(X)} = Y$.*

More precisely, for any variety $Y \subset \mathbb{G}(1, \mathbb{P}^N)$ with $1 \leq \dim Y \leq n$ and any field extension $L/K(Y)$ such that $\text{tr.deg}_k L = n$ and the linear map $\delta_{L/K(Y)}$ is zero, there exists an n -dimensional variety $X \subset \mathbb{P}^N$ such that $\overline{\gamma_X(X)} = Y$, $K(X) = L$, and the field extension $K(X)/K(Y)$ induced by γ_X coincides with the given extension $L/K(Y)$.

Remark 2.4. We give some remarks about Theorem 2.3.

- (a) For a field extension L/K , $\delta_{L/K}$ is injective if and only if L/K is separable. Hence the condition $\delta_{L/K} = 0$ is the “opposite” of the separability.
- (b) If $\dim Y = 1$, $\delta_{L/K} : \Omega_{K/k} \otimes_K L \rightarrow \Omega_{L/k}$ is zero if and only if $\delta_{L/K}$ is not injective since $\dim_L \Omega_{K/k} \otimes_K L = 1$. Thus $\delta_{L/K} = 0$ if and only if L/K is inseparable in this case by (a). Hence Theorem 2.3 generalizes Proposition 2.1.
- (c) If we replace the assumption $\delta_{L/K(Y)} = 0$ with “ $L/K(Y)$ is inseparable” as in Proposition 2.1, Theorem 2.3 does not hold in general for $\dim Y \geq 2$, even if we assume that $L/K(Y)$ is finite. See [FI2, Example 3.4].

In Theorem 2.3, we consider images and field extensions. As a geometric version of Theorem 2.3, and as a generalization of Theorem 1.5 about general fibers of Gauss maps, we have the following theorem.

Theorem 2.5. *Assume $p > 0$. Let $Y \subset \mathbb{G}(n, \mathbb{P}^N)$ be a projective variety with $\dim Y \geq 1$ and let $\mathcal{F} \subset Y \times \mathbb{P}^{N'}$ be an n -dimensional projective variety such that the first projection $f : \mathcal{F} \rightarrow Y$ is surjective. Then there exist an n -dimensional projective variety $X \subset \mathbb{P}^N$ and a generically bijective rational map $h : X \dashrightarrow \mathcal{F}$ such that the Gauss map γ_X of X is equal to $f \circ h$. In particular, $\overline{\gamma_X(X)} = Y$ holds.*

Furthermore, if we assume $n \geq N'$, we can take $X \subset \mathbb{P}^N$ such that the fiber $\overline{\gamma_X^{-1}(y)}_{\text{red}} \subset \mathbb{P}^N$ of γ_X over general $y \in Y$ is projectively equivalent to

$$F_y := f^{-1}(y)_{red} \subset \{y\} \times \mathbb{P}^{N'}.$$

$$\begin{array}{ccc} X & \xrightarrow[\text{gen. bij.}]{h} & \mathcal{F} \subset Y \times \mathbb{P}^{N'} \\ & \searrow \gamma_X & \downarrow f \\ & & Y \subset \mathbb{G}(n, \mathbb{P}^N). \end{array}$$

Roughly, Theorem 2.5 states that any surjective morphism $\mathcal{F} \rightarrow Y$ appears as a Gauss map up to generically bijective rational maps. We note that the assumption $n \geq N'$ is necessary in the last statement of Theorem 2.5 since for $y = [L] \in Y$, the general fiber $\overline{\gamma_X^{-1}(y)}_{red}$ must be contained in $L \simeq \mathbb{P}^n$.

3. IDEA OF PROOF

Let $X \subset \mathbb{P}^N$ be an n -dimensional projective variety. Then we have a rational map

$$(\gamma_X, \text{id}_X) : X \dashrightarrow \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N,$$

where id_X is the inclusion of X into \mathbb{P}^N .

Definition 3.1. We define a subvariety $\Gamma_X \subset \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N$ to be the closure of the image of (γ_X, id_X) . In other word, Γ_X is the *graph* of the rational map γ_X .

Landsberg, Piontkowski, and Furukawa characterize images of separable Gauss maps. However, it seems that such characterization does not work well for inseparable Gauss maps.

On the other hand, $(\gamma_X, \text{id}_X) : X \dashrightarrow \Gamma_X$ is birational since Γ_X is the graph of γ_X . In particular, (γ_X, id_X) is separable. Under this observation, we have the following idea:

Idea. *We characterize not images but graphs of Gauss maps to investigate not necessarily separable Gauss maps.*

To state a key theorem, we prepare some notation.

Since $\mathbb{G}(n, \mathbb{P}^N)$ is a Grassmannian, there exist locally free sheaves \mathcal{S} and \mathcal{Q} on $\mathbb{G}(n, \mathbb{P}^N)$ of ranks $N - n$ and $n + 1$ respectively with the exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow H^0(\mathbb{P}^N, \mathcal{O}(1)) \otimes \mathcal{O}_{\mathbb{G}(n, \mathbb{P}^N)} \rightarrow \mathcal{Q} \rightarrow 0.$$

Hence each point $y \in \mathbb{G}(n, \mathbb{P}^N)$ corresponds to the linear variety $\mathbb{P}(\mathcal{Q} \otimes k(y)) \subset \mathbb{P}^N$.

By the surjection $H^0(\mathbb{P}^N, \mathcal{O}(1)) \otimes \mathcal{O}_{\mathbb{G}(n, \mathbb{P}^N)} \rightarrow \mathcal{Q}$, we have a projective bundle

$$\mathcal{U} := \mathbb{P}(\mathcal{Q}) \subset \mathbb{P}(H^0(\mathbb{P}^N, \mathcal{O}(1)) \otimes \mathcal{O}_{\mathbb{G}(n, \mathbb{P}^N)}) = \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N$$

over $\mathbb{G}(n, \mathbb{P}^N)$. Hence we have the tautological invertible sheaf $\mathcal{O}_{\mathcal{U}}(1)$ on \mathcal{U} with the surjection $\text{pr}_1^* \mathcal{Q} \twoheadrightarrow \mathcal{O}_{\mathcal{U}}(1)$. By definition, it holds that

$$\mathcal{U} = \{([L], x) \in \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N \mid x \in L\},$$

that is, \mathcal{U} is the incidence variety.

For a variety $X' \subset \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N$, let $\text{pr}_1 : X' \rightarrow \mathbb{G}(n, \mathbb{P}^N)$ and $\text{pr}_2 : X' \rightarrow \mathbb{P}^N$ be the first and second projections respectively. Since $\Omega_{\mathbb{G}(n, \mathbb{P}^N)} = \mathcal{Q}^\vee \otimes \mathcal{S}$ holds, there exists a natural homomorphism

$$\text{pr}_1^*(\mathcal{Q}^\vee \otimes \mathcal{S}) = \text{pr}_1^* \Omega_{\mathbb{G}(n, \mathbb{P}^N)} \rightarrow \Omega_{X'}$$

for a variety $X' \subset \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N$. This homomorphism induces

$$\Phi : \text{pr}_1^* \mathcal{Q}^\vee \rightarrow \Omega_{X'} \otimes \text{pr}_1^* \mathcal{S}^\vee.$$

For an n -dimensional projective variety $X \subset \mathbb{P}^N$, the graph Γ_X is contained in \mathcal{U} since $x \in \gamma_X(x) = \mathbb{T}_x X$ for a smooth point $x \in X$. Hence we have the following commutative diagram:

$$\begin{array}{ccc} & & \Gamma_X \subset \mathcal{U} \subset \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^N \\ & \text{pr}_2 \nearrow & \downarrow \text{pr}_1 \\ X & \xrightarrow{\gamma_X} & \mathbb{G}(n, \mathbb{P}^N) \end{array}$$

(dashed arrow from X to Γ_X is labeled (γ_X, id_X))

The following theorem gives a characterization of graphs of Gauss maps. We note that this theorem holds for any characteristic.

Theorem 3.2. *Let $X' \subset \mathcal{U}$ be a subvariety of dimension n such that $\text{pr}_2 : X' \rightarrow \mathbb{P}^N$ is generically finite and separable. Then the following are equivalent:*

- (i) $X' = \Gamma_X$ holds for some n -dimensional variety $X \subset \mathbb{P}^N$.
- (ii) $X' = \Gamma_{X_0}$ holds for $X_0 = \text{pr}_2(X')$.
- (iii) The composite homomorphism $\mathcal{O}_{\mathcal{U}}(-1)|_{X'} \hookrightarrow \text{pr}_1^* \mathcal{Q}^\vee \xrightarrow{\Phi} \Omega_{X'} \otimes \text{pr}_1^* \mathcal{S}^\vee$ is the zero map at the generic point of X' .

Proof. See [FI2]. □

Remark 3.3. By definition, Φ is the zero map at the generic point of X' if and only if so is $\text{pr}_1^*(\mathcal{Q}^\vee \otimes \mathcal{S}) = \text{pr}_1^* \Omega_{\mathbb{G}(n, \mathbb{P}^N)} \rightarrow \Omega_{X'}$, i.e. the $K(X)$ -linear map $\delta_{K(X')/K(\text{pr}_1(X'))}$ is zero. Hence (iii) in Theorem 3.2 holds automatically if $\delta_{K(X')/K(\text{pr}_1(X'))} = 0$.

By using Theorem 3.2, we can show Theorem 2.3 as follows.

Sketch of proof of Theorem 2.3. Assume that there exists an n -dimensional variety $X' \subset \mathcal{U}$ which satisfies

- $\text{pr}_2 : X' \rightarrow \mathbb{P}^N$ is generically finite and separable,
- $Y = \text{pr}_1(X')$,
- the field extension $K(X')/K(Y)$ induced by pr_1 coincides with the given extension $L/K(Y)$.

Then X' is the graph of γ_X for $X = \text{pr}_2(X') \subset \mathbb{P}^N$ by Theorem 3.2 and the assumption $\delta_{L/K(Y)} = 0$ (see Remark 3.3), and this X satisfies the conditions in Theorem 2.3.

Hence it suffices to find such X' . To find such X' , it is enough to find $x_1, \dots, x_n \in L$ such that

- $L/k(x_1, \dots, x_n)$ is finite and separable,
- $L = K(Y)(x_1, \dots, x_n)$.

In fact, we take X' to be the image of

$$\text{Spec } L \xrightarrow{(f,g)} Y \times \mathbb{P}^n \subset \mathbb{G}(n, \mathbb{P}^N) \times \mathbb{P}^n \xrightarrow{\text{bir}} \mathbb{P}(\mathcal{Q}) = \mathcal{U},$$

where $f : \text{Spec } L \rightarrow Y$ is induced by $L/K(Y)$ and $g : \text{Spec } L \rightarrow \mathbb{P}^n$ is the morphism which is parametrized as $[1 : x_1 : \dots : x_n]$.

By a suitable argument about the generators of field extensions, we can find such x_i and Theorem 2.3 is proved. \square

In Theorem 2.3, we consider only field extensions $L/K(Y)$ with $\delta_{L/K(Y)} = 0$. How about field extensions with $\delta_{L/K(Y)} \neq 0$? By using Theorem 3.2, we can show the following theorem about inseparable field extensions L/K , not necessarily $\delta_{L/K} = 0$.

Theorem 3.4. *Let L/K be an inseparable field extension. If $p \geq 3$ or $\text{rank}_L \delta_{L/K}$ is even, there exists a hypersurface $X \subset \mathbb{P}^{n+1}$ for $n = \text{tr.deg}_k L$ such that the extension $K(X)/K(\gamma_X(X))$ induced by γ_X coincides with the given extension L/K .*

The difference of Theorems 2.3 and 3.4 are as follows: In Theorem 2.3, we fix an embedding $\text{Spec } K \hookrightarrow \mathbb{G}(n, \mathbb{P}^N)$ as $Y \subset \mathbb{G}(n, \mathbb{P}^N)$, and consider field extensions L/K such that $\delta_{L/K}$ is zero. In Theorem 3.4, we do not fix an embedding $\text{Spec } K \hookrightarrow \mathbb{G}(n, \mathbb{P}^N)$, and consider L/K such that $\delta_{L/K}$ is not necessarily zero.

In the proof of Theorem 2.3, the condition (iii) in Theorem 3.2 automatically holds since $\Phi = 0$, which follows from the assumption $\delta_{L/K} = 0$. On the other hand, Φ is not zero in the setting of Theorem 3.4. Hence we need to take an embedding $\text{Spec } K \hookrightarrow \mathbb{G}(n, \mathbb{P}^{n+1})$ and $x_1, \dots, x_n \in L$ carefully so that the condition (iii) in Theorem 3.2 holds.

Remark 3.5. If $p = 2$, the behavior of Gauss maps is sometimes different from that in other characteristics. For example,

- For any hypersurface $X \subset \mathbb{P}^{n+1}$, $\text{rank}_{K(X)} \delta_{K(X)/K(\gamma(X))}$ is even. Hence the statement of Theorem 3.4 does not hold if $p = 2$ and $\text{rank}_L \delta_{L/K}$ is odd.
- For any variety $X \subset \mathbb{P}^N$ (which is not necessarily a hypersurface), $\text{rank}_{K(X)} \delta_{K(X)/K(\gamma(X))}$ cannot be equal to one.

By Theorem 3.4 and Remark 3.5, we have the following question:

Question 3.6. *Assume $p = 2$. Let L/K be an inseparable field extension with $n = \text{tr.deg}_k L$ and $\text{rank}_L \delta_{L/K}$ is odd. Then is there an n -dimensional variety $X \subset \mathbb{P}^N$ with $N \geq n + 2$ such that $K(X)/K(\gamma(X)) = L/K$?*

Unfortunately, we do not know the answer of this question.

REFERENCES

- [FP] G. Fischer and J. Piontowski, *Ruled varieties*, Advanced Lectures in Mathematics, Friedr. Vieweg & Sohn, Braunschweig, 2001.
- [Fuk1] S. Fukasawa, Developable varieties in positive characteristic. *Hiroshima Math. J.* **35** (2005), 167–182.
- [Fuk2] ———, Varieties with non-linear Gauss fibers. *Math. Ann.* **334** (2006), 235–239.
- [Fur] K. Furukawa, Duality with expanding maps and shrinking maps, and its applications to Gauss maps, *Math. Ann.* **358** (2014), 403–432.
- [FI1] K. Furukawa and A. Ito, Gauss maps of toric varieties, arXiv:1403.0793.
- [FI2] ———, On Gauss maps in positive characteristics in view of images, fibers, and field extensions, in arXiv.
- [GH] P. Griffiths and J. Harris, Algebraic geometry and local differential geometry. *Ann. Sci. Ecole Norm. Sup. (4)* **12** (1979), 355–432.
- [IL] T. A. Ivey and J. M. Landsberg, *Cartan for beginners: differential geometry via moving frames and exterior differential systems*. Grad. Stud. Math. **61**, Amer. Math. Soc., Providence, 2003.
- [Ka1] ———, On the Gauss maps of space curves in characteristic p . *Compositio Math.* **70** (1989), 177–197.
- [Ka2] H. Kaji, On the space curves with the same image under the Gauss maps. *Manuscripta Math.* **80** (1993), 249–258.
- [Ra] J. Rathmann, The uniform position principle for curves in characteristic p . *Math. Ann.* **276** (1987), 565–579.
- [Za] F. L. Zak, *Tangents and secants of algebraic varieties*. Transl. Math. Monographs **127**, Amer. Math. Soc., Providence, 1993.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN
E-mail address: aito@math.kyoto-u.ac.jp